arXiv:0904.1261v1 [math.AP] 8 Apr 2009

EXISTENCE OF A DEGENERATE SINGULARITY IN THE HIGH ACTIVATION ENERGY LIMIT OF A REACTION-DIFFUSION EQUATION

GEORG S. WEISS AND GUANGHUI ZHANG

Abstract. We consider the singular perturbation problem

$$\Delta u_\epsilon = \beta_\epsilon(u_\epsilon),$$

where $\beta_\epsilon(s) = \frac{1}{\epsilon}\beta(\frac{s}{\epsilon})$, β is a Lipschitz continuous function such that $\beta > 0$ in $(0,1)$, $\beta \equiv 0$ outside $(0,1)$ and $\int_0^1 \beta(s)\, ds = \frac{1}{2}$.

We construct an example exhibiting a *degenerate singularity* as $\epsilon_k \searrow 0$. More precisely, there is a sequence of solutions $u_{\epsilon_k} \to u$ as $k \to \infty$, and there exists $x^0 \in \partial\{u > 0\}$ such that

$$\frac{u(x^0 + r\cdot)}{r} \to 0 \text{ as } r \to 0.$$

Known results suggest that this singularity must be *unstable*, which makes it hard to capture analytically and numerically. Our result answers a question raised by Jean-Michel Roquejoffre at the FBP'08 in Stockholm.

1. Introduction

In this paper we are concerned with the singular perturbation problem

$$\Delta u_\epsilon = \beta_\epsilon(u_\epsilon) \text{ in } \Omega, \tag{1.1}$$

where Ω is a bounded domain in $\mathbb{R}^n$, $\epsilon > 0$ and $\beta_\epsilon(s) = \frac{1}{\epsilon}\beta(\frac{s}{\epsilon})$. Here β is a Lipschitz continuous function such that $\beta > 0$ in $(0,1)$, $\beta \equiv 0$ outside $(0,1)$ and $\int_0^1 \beta(s)\, ds = \frac{1}{2}$.

2000 *Mathematics Subject Classification.* Primary 35R35, Secondary 35J60.

Key words and phrases. Free boundary, singular point, unstable singularity.

G.S. Weiss has been partially supported by the Grant-in-Aid 18740086 of the Japanese Ministry of Education, Culture, Sports, Science and Technology.

This problem arises in the mathematical analysis of equidiffusional flames (see [3], [4]), in which case ϵ is proportional to the inverse of the activation energy.

Formally, as $\epsilon \to 0$, the solutions u_ϵ converge to a solution u of the free boundary problem

$$\begin{array}{lll} & \Delta u = 0 & \text{in } \Omega \setminus \partial\{u > 0\}, \\ (1.2) & & \\ & u = 0,\ (u_\nu^+)^2 - (u_\nu^-)^2 = 1 & \text{on } \partial\{u > 0\}. \end{array}$$

On a rigorous level, L. Caffarelli established a locally uniformly Lipschitz estimate for bounded solutions $\{u_\epsilon\}$ (see [5]). In [10], C. Lederman and N. Wolanski proved that u is a viscosity solution of (1.2). They also proved that u satisfies the free boundary condition in a pointwise sense at non-degenerate free boundary points at which there is an inward unit normal of $\{u > 0\}$ in the measure theoretic sense. Here "non-degenerate" means that

$$r^{-n-1} \int_{B_r(x)} u^+ \geq c > 0 \text{ for } r \leq r_0.$$

A related convergence result has been proved for the p-Laplace operator in [6].

From [1, Lemma 3.4] we know that all free boundary points of local minimizers are non-degenerate. This suggests that if there are degenerate free boundary points of a limit function u, then they must be unstable.

It has been known that cross-shaped free boundaries occur for domain variation solutions (see the Introduction of [12]), for example

$$u(x_1, x_2) = |x_1^2 - x_2^2| \text{ (cf. Figure 1).}$$

John Andersson suggested a triple junction example (see Figure 2) where the solution is homogeneous of degree 3/2; domain variation solutions with a degree of homogeneity $\in (1, 3/2)$ are not possible. However it is not so obvious whether these examples arise as limits of the reaction-diffusion equation (1.1). This is the question Jean-Michel Roquejoffre posed at the FBP'08 in Stockholm. The main result of the present paper gives a partial answer to that question:



Figure 1. Cross-like Singularity



Figure 2. Triple Junction Singularity

Theorem A. *There exist uniformly bounded solutions* $u_{\epsilon_k} : B_1(0) \subset \mathbb{R}^2 \to \mathbb{R}$ *such that* $\epsilon_k \to 0, u_{\epsilon_k} \to u$ *as* $k \to \infty$, *and there is a degenerate free boundary point of* u *in* $B_1(0)$. *More precisely, there is* $x^0 \in B_1(0) \cap \partial\{u > 0\}$ *such that*

$$\lim_{r \to 0} \frac{u(x^0 + rx)}{r} = 0 \text{ for every } x \in \mathbb{R}^2.$$

Remark 6.5 shows that we may construct degenerate points with an arbitrarily high number of symmetry lines. See Figure 3 for an example with 4 symmetry lines.
Section 3 also suggests a *numerical* approach for the construction of this unstable solution which would be otherwise hard to capture.

2. Notation

Throughout this article $\mathbb{R}^n$ will be equipped with the Euclidean inner product $x \cdot y$ and $B_r(x^0)$ will denote the open ball of center x^0 with



FIGURE 3. Imaginable ϵ-solution

radius r. When not specified, x^0 is assumed to be 0. We will use the polar coordinates (r, θ) in $\mathbb{R}^2$. We use the n-dimensional Lebesgue measure $\mathcal{L}^n$ and the $m-$dimensional Hausdorff measure $\mathcal{H}^m$. We shall often use abbreviations for inverse images like $\{u > 0\} := \{x \in \Omega : u(x) > 0\}$.

3. CONSTRUCTION OF $\epsilon-$SOLUTIONS

We are going to use a method introduced in [2].

Let $K = \{(r\cos\theta, r\sin\theta) : 0 < r < 1, 0 < \theta < \pi/m\}$, where m is an integer ≥ 3. For $g \in C^{\alpha_0}(\partial B_1 \cap \partial K)$, $\alpha_0 \geq \alpha > 0$, $C_g^\alpha(\bar{K})$ will denote the subspace of $C^\alpha(\bar{K})$ consisting of all functions with boundary values g on $\partial B_1 \cap \partial K$. We consider the following problem:

$$\begin{aligned} &\Delta u(x) = \beta_\epsilon(u(x) - u(0) + \epsilon) && \text{in } K, \\ &u = g && \text{on } \partial K \cap \partial B_1, \\ &\tfrac{\partial u}{\partial \nu} = 0 && \text{on } \partial K \setminus \partial B_1. \end{aligned} \tag{3.1}$$

Let $T = T_{\epsilon,g} : C_g^\alpha(\bar{K}) \to C_g^\alpha(\bar{K})$ be the operator defined by

$$\begin{aligned} &\Delta T(u) = \beta_\epsilon(u(x) - u(0) + \epsilon) && \text{in } K, \\ &T(u) = g && \text{on } \partial K \cap \partial B_1, \\ &\tfrac{\partial T(u)}{\partial \nu} = 0 && \text{on } \partial K \setminus \partial B_1. \end{aligned}$$

Using Schauder's fixed point theorem, we can prove the existence of a solution of equation (3.1):

Proposition 3.1. *For each $g \in C^{\alpha_0}(\partial B_1 \cap \partial K)$, T has a fixed point $u_\epsilon \in C^\alpha_g(\bar{K})$ for some $\alpha > 0$ depending only on α_0.*

Proof. Let $v \in C^\alpha_g(\bar{K})$, $F(x) = \beta_\epsilon(v(x) - v(0) + \epsilon)$. Then there is a $W^{1,2}(K)$-solution u of the problem:

$$\begin{aligned} &\Delta u = F && \text{in } K,\\ &u = g && \text{on } \partial K \cap \partial B_1,\\ &\tfrac{\partial u}{\partial \nu} = 0 && \text{on } \partial K \setminus \partial B_1. \end{aligned}$$

We extend u to a function $\tilde{u} : B_1 \setminus \{0\} \to \mathbb{R}$ by even reflection:

$$\begin{aligned} &\tilde{u}(r, \pi/m + \theta) = u(r, \pi/m - \theta), && \text{if } 0 \le \theta \le \pi/m,\\ &\tilde{u}(r, j\pi/m + \theta) = \tilde{u}(r, j\pi/m - \theta), && \text{if } 0 \le \theta \le \pi/m, \end{aligned}$$

for $j = 1, 2, \dots, 2m-1$. It follows that $\tilde{u}$ is a solution of

$$\begin{aligned} &\Delta \tilde{u} = \tilde{F} && \text{in } B_1 \setminus \{0\},\\ &\tilde{u} = \tilde{g} && \text{on } \partial B_1, \end{aligned}$$

where $\tilde{F}$ and $\tilde{g}$ are defined by even reflection. As the origin is a set of vanishing 2-capacity (see [7]), $\tilde{u}$ is a weak solution of $\Delta \tilde{u} = \tilde{F}$ in B_1.

Applying the regularity theory of elliptic equations (see for example [8, Corollary 9.29]), we see that $\tilde{u} \in C^\alpha(\bar{B}_1)$ for small α, and that $||\tilde{u}||_{C^{0,\alpha}(B_1)} \le C$, where $\alpha \in (0, \alpha_0)$ and C are constants depending on α_0, $||g||_{C^{0,\alpha_0}}$ and $||F||_{L^\infty(K)}$. Hence T is a continuous compact linear operator from $C^\alpha_g(\bar{K})$ into itself for small $\alpha > 0$, and

$$||T_{\epsilon,g}(w)||_{C^\alpha(\bar{K})} \le C$$

for every $\sigma \in [0, 1]$ and every solution of the equation

$$\begin{aligned} &\Delta w = \sigma \beta_\epsilon(w(x) - w(0) + \epsilon) && \text{in } K,\\ &w = g && \text{on } \partial K \cap \partial B_1,\\ &\tfrac{\partial w}{\partial \nu} = 0 && \text{on } \partial K \setminus \partial B_1; \end{aligned}$$

here C is a constant depending only on g, β and ϵ.

From Schauder's fixed point theorem (see [8, Chapter 11]) we infer that $T_{\epsilon,g}$ has a fixed point $u_\epsilon \in C^\alpha_g(\bar{K})$. □

4. CONVERGENCE

From now on all of our functions are defined in $B_1(0)$ by the above reflection. We choose one non-constant $g \in C^2(\bar{B})$.

Let $v_\epsilon(x) = u_\epsilon(x) - u_\epsilon(0) + \epsilon$. Then v_ϵ is a solution of

$$\begin{aligned} &\Delta v_\epsilon(x) = \beta_\epsilon(v_\epsilon(x)) && \text{in } B_1,\\ &v_\epsilon(0) = \epsilon, \\ &v_\epsilon = g - u_\epsilon(0) + \epsilon && \text{on } \partial B_1. \end{aligned}$$

By the maximum principle, $|u_\epsilon(0)| \le ||g||_{L^\infty(B_1)}$, therefore $||v_\epsilon||_{L^\infty(\bar{B}_1)} \le C$, where C depends only on $||g||_{L^\infty(B_1)}$.

First we state a result from [5] proving a uniform Lipschitz estimate.

Proposition 4.1. *Let w_ϵ be a family of solutions to $\Delta w_\epsilon = \beta_\epsilon(w_\epsilon)$ in a domain $\Omega \subset \mathbb{R}^n$ such that $||w_\epsilon||_{L^\infty(\Omega)} \le C$ for some $C < \infty$. Let $K \subset \Omega$ be a compact set and let $\tau > 0$ be such that $B_\tau(x^0) \subset \Omega$ for every $x^0 \in K$. Then there exists a constant $L = L(\tau, C)$, such that*

$$|\nabla w_\epsilon(x)| \le L \quad \text{for } x \in K.$$

If the boundary data is smooth enough, we have the following gradient estimate near the boundary (see [9]).

Proposition 4.2. *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with smooth boundary and let $w_\epsilon \in C^2(\Omega) \cap C^0(\bar{\Omega})$ be a solution of*

$$\begin{aligned} &\Delta w_\epsilon(x) = \beta_\epsilon(w_\epsilon(x)) && \text{in } \Omega,\\ &w_\epsilon = f && \text{on } \partial\Omega, \end{aligned}$$

where $f \in C^2(\bar{\Omega})$, $||f||_{C^2(\bar{\Omega})} < C_1$. Then there exist $\epsilon_0 > 0, C < \infty$ such that

$$|\nabla w_\epsilon(x)| \le C(1 + |\log d(x, \partial\Omega)|)$$

for $\epsilon \in (0, \epsilon_0)$, $x \in \Omega$, where $d(x, \partial\Omega)$ is the distance from x to the boundary of Ω.

Now we are ready to prove the convergence of $\{v_{\epsilon_i}\}$.

Proposition 4.3. *There exist a sequence $\{v_{\epsilon_i}\}$ and a locally Lipschitz continuous function v_0 such that*
1) $v_{\epsilon_i} \to v_0$ uniformly on $\bar{B}_1$,

2) $v_{\epsilon_i} \to v_0$ *in* $W^{1,2}_{\text{loc}}(B_1)$,
3) v_0 *is harmonic in* $B_1 \setminus \partial\{v_0 > 0\}$,
4) $\Delta v_{\epsilon_i} \to \mu$ *as measures on* B_1*; here* μ *is a locally finite non-negative measure supported on the free boundary* $\partial\{v_0 > 0\}$*. Therefore*

$$\Delta v_0 = \mu \qquad \text{in } B_1,$$

that is

$$\int_{B_1} \nabla v_0 \cdot \nabla \phi \; dx + \int_{B_1} \phi \; d\mu = 0$$

for every $\phi \in C^1_0(B_1)$.
5) $v_0(r, j\pi/m+\theta) = v_0(r, j\pi/m-\theta)$ *for* $0 \le \theta \le \pi/m$ *and* $1 \le j \le 2m$.

Proof. 1) By Proposition 4.2, $\{v_\epsilon\}$ is equicontinuous. Since $\{v_\epsilon\}$ is uniformly bounded in $C^0(\bar{B}_1)$, by the Arzela-Ascoli theorem there is a sequence $\{v_{\epsilon_i}\}$ that converges to some v_0 in $C^0(\bar{B}_1)$. By Proposition 4.1, it is easy to see that v_0 is locally Lipschitz continuous.

2) For $0 \le \eta \in C^\infty_0(B_1)$, we obtain from the local Lipschitz bound that

$$\int_{B_1} \eta |\nabla v_0|^2 \; dx \le \liminf_{\epsilon_i \to 0} \int_{B_1} \eta |\nabla v_{\epsilon_i}|^2 \; dx. \tag{4.1}$$

If we prove the converse inequality, the result will follow. Multiplying the ϵ_i-equation (1.1) by ηv_{ϵ_i}, we obtain that

$$\int_{B_1} v_{\epsilon_i} \nabla \eta \cdot \nabla v_{\epsilon_i} \; dx + \int_{B_1} \eta |\nabla v_{\epsilon_i}|^2 \; dx = -\int_{B_1} \eta v_{\epsilon_i} \beta_{\epsilon_i}(v_{\epsilon_i}) \; dx \le 0.$$

Letting $\epsilon_i \to 0$, we get

$$\limsup_{\epsilon_i \to 0} \int_{B_1} \eta |\nabla v_{\epsilon_i}|^2 \; dx \le -\int_{B_1} v_0 \nabla \eta \cdot \nabla v_0 \; dx. \tag{4.2}$$

For each $\delta > 0$, we define

$$v_\delta(x) = \begin{cases} v_0(x) - \delta, & \text{if } v_0(x) > \delta, \\ v_0(x) + \delta, & \text{if } v_0(x) < -\delta, \\ 0 & \text{otherwise.} \end{cases}$$

Multiplying the ϵ_i-equation by ηv_δ and integrating, we have

$$\int_{B_1} v_\delta \nabla\eta \cdot \nabla v_{\epsilon_i}\, dx + \int_{B_1} \eta \nabla v_\delta \cdot \nabla v_{\epsilon_i}\, dx = 0.$$

Letting $\epsilon_i \to 0$, it follows that

$$\int_{B_1} v_\delta \nabla\eta \cdot \nabla v_0\, dx + \int_{B_1} \eta |\nabla v_\delta|^2\, dx = 0,$$

and letting $\delta \to 0$, we get

$$\int_{B_1} v_0 \nabla\eta \cdot \nabla v_0\, dx + \int_{B_1} \eta |\nabla v_0|^2\, dx = 0.$$

Using (4.2), we obtain

$$\limsup_{\epsilon_i \to 0} \int_{B_1} \eta |\nabla v_{\epsilon_i}|^2\, dx \le \int_{B_1} \eta |\nabla v_0|^2\, dx. \tag{4.3}$$

Combining (4.1) and (4.3), we obtain

$$\lim_{\epsilon_i \to 0} \int_{B_1} \eta |\nabla v_{\epsilon_i}|^2\, dx = \int_{B_1} \eta |\nabla v_0|^2\, dx, \tag{4.4}$$

which implies that

$$\eta^{1/2} \nabla v_{\epsilon_i} \to \eta^{1/2} \nabla v_0 \ \text{ in } L^2(B_1).$$

For η such that $\eta = 1$ in $D \subset\subset \Omega$, it follows that

$$\nabla v_{\epsilon_i} \to \nabla v_0 \ \text{ in } L^2(D).$$

Therefore $v_\epsilon \to v_0$ strongly in $W^{1,2}(D)$.

By a diagonal sequence argument, assertion 2) of the proposition follows.

3) Since v_0 is continuous, the sets $\{v_0 > 0\}$ and $\{v_0 < 0\}$ are open. Let $x^0 \in \{v_0 > 0\}$. From the fact that $v_{\epsilon_i} \to v_0$ uniformly on B_1, there exist $r > 0$ and $N \in \mathbb{N}$ such that $v_{\epsilon_i}(x) \ge v_0(x^0)/2 > 0$ for $x \in B_r(x^0)$, $i \ge N$. Thus v_{ϵ_i} is harmonic in $B_r(x^0)$ for $i \ge N$ and the same fact holds for v_0. In the same way we prove that v_0 is superharmonic in $\{v_0 \le 0\}^0$. On the other hand, v_{ϵ_i} is subharmonic in B_1, so that v_0 is also subharmonic. Therefore v_0 is harmonic in $\{v_0 \le 0\}^0$ and 3) follows.

4) Multiplying equation $\Delta v_{\epsilon_i} = \beta_{\epsilon_i}(v_{\epsilon_i})$ by $\phi \in C_0^\infty(B_1)$ and integrating by parts, we get

$$\int_{B_1} \nabla v_{\epsilon_i} \cdot \nabla \phi \, dx + \int_{B_1} \beta_{\epsilon_i}(v_{\epsilon_i})\phi \, dx = 0.$$

We know that $v_{\epsilon_i} \to v_0$ in $W^{1,2}_{\mathrm{loc}}(B_1)$, hence

$$\int_{B_1} \nabla v_{\epsilon_i} \cdot \nabla \phi \, dx \to \int_{B_1} \nabla v_0 \cdot \nabla \phi \, dx. \tag{4.5}$$

On the other hand,

$$\int_{B_1} \beta_{\epsilon_i}(v_{\epsilon_i})\phi \, dx \le C.$$

This L^1-bound implies that there exists a locally finite non-negative measure μ such that for a subsequence which we still call $\{v_{\epsilon_i}\}$, $\beta_{\epsilon_i}(v_{\epsilon_i}) \to \mu$ as measures in B_1. Passing to the limit in (4.5), we get

$$\int_{B_1} \nabla v_0 \cdot \nabla \phi \, dx + \int_{B_1} \phi \, d\mu = 0,$$

which implies

$$\Delta v_0 = \mu \qquad \text{in } B_1.$$

Since we know that v_0 is harmonic in $B_1 \setminus \partial\{v_0 > 0\}$, we conclude that $\operatorname{supp} \mu \subset \partial\{v_0 > 0\}$.

5) follows from the fact that v_{ϵ_i} has been defined via even reflection. This property is preserved by the uniform limit v_0. □

Let $\mathcal{B}_\epsilon(z) = \int_0^z \beta_\epsilon(s)\, ds$ and $\chi_\epsilon(x) = 2\mathcal{B}_\epsilon(v_\epsilon(x))$. It follows that $0 \le \chi_\epsilon(x) \le 1$, and the uniformly Lipschitz estimate for v_ϵ implies the relative compactness of χ_ϵ in $L^1_{\mathrm{loc}}(B_1)$:

Proposition 4.4. *$\{\chi_\epsilon\}$ is precompact in $L^1(D)$ for each $D \subset\subset B_1$.*

Proof. By Proposition 4.1, there exists an $L > 0$ such that $||\nabla v_\epsilon||_{L^\infty(D)} \le L$, hence

$$\int_D |\nabla \chi_\epsilon| \, dx = 2\int_D |\beta_\epsilon(v_\epsilon)\nabla v_\epsilon| \, dx \le 2L \int_D \beta_\epsilon(v_\epsilon)\, dx \le C.$$

Therefore $\{\chi_\epsilon\}$ is bounded in $W^{1,1}(D)$. By the Sobolev embedding theorem $\{\chi_\epsilon\}$ is precompact in $L^1(D)$. □

Thus we may assume that $\chi_{\epsilon_i}(x) \to \chi_0(x)$ locally in L^1. Similar to [12, Lemma 4.1], we have

Proposition 4.5. *$\chi_0(x) \in \{0,1\}$ for a.e. $x \in B_1$.*

Proof. For $\delta \in (0,1/4)$, $\mathcal{B}(s) = 2\int_0^s \beta(t)\,dt$ and $A_\delta = \{x \in D \subset\subset B_1 : \chi_0 \in [2\delta, 1-2\delta]\}$ we have

$$\begin{aligned}\mathcal{L}^2(A_\delta) &\le \mathcal{L}^2(A_\delta \cap \{\chi_{\epsilon_m} \notin [\delta, 1-\delta]\}) + \mathcal{L}^2(A_\delta \cap \{\mathcal{B}(\frac{v_{\epsilon_m}}{\epsilon_m}) \in [\delta, 1-\delta]\})\\ &\le \mathcal{L}^2(D \cap \{|\chi_{\epsilon_m} - \chi_0| > \delta\}) + \mathcal{L}^2(D \cap \{\frac{v_{\epsilon_m}}{\epsilon_m} \in [\mathcal{B}^{-1}(\delta), \mathcal{B}^{-1}(1-\delta)]\})\\ &\le \mathcal{L}^2(D \cap \{|\chi_{\epsilon_m} - \chi_0| > \delta\}) + C_1(\delta,\beta)\epsilon_m \int_D \beta_{\epsilon_m}(v_{\epsilon_m})\,dx\\ &\le \mathcal{L}^2(D \cap \{|\chi_{\epsilon_m} - \chi_0| > \delta\}) + C_2(\delta,\beta)\epsilon_m \to 0\end{aligned}$$

as $m \to \infty$. □

5. Monotonicity formula

In this section we state a monotonicity formula proved in [13], which is a key tool in constructing a degenerate singularity.

Proposition 5.1. *Let $x^0 \in B_1$, and*

$$\Phi_\epsilon(r) = r^{-2}\int_{B_r(x^0)} |\nabla v_\epsilon|^2 + r^{-2}\int_{B_r(x^0)} \chi_\epsilon - r^{-3}\int_{\partial B_r(x^0)} v_\epsilon^2\,d\mathcal{H}^1.$$

Then Φ_ϵ satisfies the monotonicity formula

$$\Phi_\epsilon(\sigma) - \Phi_\epsilon(\rho) \ge \int_\rho^\sigma r^{-2}\int_{\partial B_r(x^0)} 2(\nabla v_\epsilon \cdot \nu - \frac{v_\epsilon(x)}{r})^2\,d\mathcal{H}^1 dr \quad \text{for } 0 < \rho < \sigma.$$

Letting $\epsilon \to 0$,

$$\Phi(r) = r^{-2}\int_{B_r(x^0)} |\nabla v_0|^2 + r^{-2}\int_{B_r(x^0)} \chi_0 - r^{-3}\int_{\partial B_r(x^0)} {v_0}^2\,d\mathcal{H}^1$$

satisfies the monotonicity formula

$$\Phi(\sigma) - \Phi(\rho) \ge \int_\rho^\sigma r^{-2}\int_{\partial B_r(x^0)} 2(\nabla v_0 \cdot \nu - \frac{v_0(x)}{r})^2\,d\mathcal{H}^1 dr \quad \text{for } 0 < \rho < \sigma.$$

6. Existence of a Degenerate Free Boundary Point

In this section we are going to prove the existence of a degenerate singular free boundary point for v_0. We start with some lemmas.

Lemma 6.1. *Let w be a solution of*

$$\Delta w = \beta_\epsilon(w) \text{ in } \Omega \subset \mathbb{R}^n \text{ and let } \chi_\epsilon(x) = 2\mathcal{B}_\epsilon(w(x)).$$

Then

$$\int_\Omega |\nabla w|^2 \text{div } \phi - 2\nabla w D\phi \nabla w + \chi_\epsilon \text{div } \phi \, dx = 0$$

for $\phi \in C_0^1(\Omega; \mathbb{R}^n)$.

Proof. Integrating by parts, we get

$$\begin{aligned}
&\int_\Omega |\nabla w|^2 \text{div } \phi - 2\nabla w D\phi \nabla w + \chi_\epsilon \text{div } \phi \, dx \\
&= \int_\Omega \sum_{i=1}^n \left(\frac{\partial w}{\partial x_i}\right)^2 \sum_{j=1}^n \frac{\partial \phi_j}{\partial x_j} - 2 \sum_{i,j=1}^n \frac{\partial w}{\partial x_i} \frac{\partial \phi_j}{\partial x_i} \frac{\partial w}{\partial x_j} + \chi_\epsilon \text{div } \phi \, dx \\
&= -2 \int_\Omega \sum_{i,j=1}^n \phi_j \frac{\partial w}{\partial x_i} \frac{\partial^2 w}{\partial x_i \partial x_j} - \sum_{i,j=1}^n \phi_j \frac{\partial}{\partial x_i} \left(\frac{\partial w}{\partial x_i} \frac{\partial w}{\partial x_j} \right) + \beta_\epsilon(w) \nabla w \cdot \phi \, dx \\
&= 2 \int_\Omega \sum_{i,j=1}^n \phi_j \frac{\partial w}{\partial x_j} \frac{\partial^2 w}{\partial x_i^2} - \beta_\epsilon(w) \nabla w \cdot \phi \, dx \\
&= 2 \int_\Omega \nabla w \cdot \phi (\Delta w - \beta_\epsilon(w)) \, dx \\
&= 0.
\end{aligned}$$

□

Lemma 6.2. *If $v_0 \equiv 0$ in B_r for some $r > 0$, then $\chi_0 \equiv 1$ in B_r.*

Proof. By Lemma 6.1 we have

$$\int_{B_1} |\nabla v_{\epsilon_i}|^2 \text{div } \phi - 2\nabla v_{\epsilon_i} D\phi \nabla v_{\epsilon_i} + \chi_{\epsilon_i} \text{div } \phi \, dx = 0$$



FIGURE 4. The set P_i

for $\phi \in C^\infty_0(B_r;\mathbb{R}^2)$. Letting $\epsilon_i \to 0$, we obtain from the strong convergence of v_{ϵ_i} that

$$\begin{aligned} 0 &= \int_{B_1} |\nabla v_0|^2 \text{div } \phi - 2\nabla v_0 D\phi \nabla v_0 + \chi_0 \text{div } \phi \, dx \\ &= \int_{B_r} \chi_0 \text{div } \phi \, dx, \end{aligned}$$

hence $\chi_0 \equiv$ const in B_r. We know that $\chi_0(x) \in \{0,1\}$ for a.e. $x \in B_1$, therefore $\chi_0(x) \equiv 0$ or $\chi_0(x) \equiv 1$ in B_r.

For $0 < c \le 1/4$, let $P_i = \{x \in B_r : c\epsilon_i \le v_{\epsilon_i}(x) \le (1-c)\epsilon_i\}$ and $Q_i = \{x \in B_r : v_{\epsilon_i}(x) > c\epsilon_i\}$. Supposing towards a contradiction that $\chi_0 \equiv 0$ in B_r, and recalling that $\chi_{\epsilon_i} \to \chi_0$ in $L^1(B_r)$ we have

$$\mathcal{L}^2(Q_i) \le \mathcal{L}^2(\{x : \chi_{\epsilon_i} > 2\mathcal{B}(c)\}) \to 0$$

as $i \to \infty$.

Let $\delta \in (0, r/4)$, $I_{\epsilon_i} = \{t : \delta \le t \le r, \min_{x\in\partial B_t} v_{\epsilon_i}(x) \ge \epsilon_i/2\}$ and $I^c_{\epsilon_i} = [\delta, r] \setminus I_{\epsilon_i}$. Then we have $\mathcal{L}^1(I_{\epsilon_i}) \le \mathcal{L}^2(Q_i)/\delta \to 0$ as $i \to \infty$, hence there exists an $\epsilon_0 > 0$ such that $\mathcal{L}^1(I^c_{\epsilon_i}) > r/2$ for $\epsilon_i < \epsilon_0$. As v_{ϵ_i} is subharmonic in B_1, $\epsilon_i = v_{\epsilon_i}(0) \le \max_{\partial B_t} v_{\epsilon_i}$ for every $t \in [0, r)$. It follows that for each $t \in I^c_{\epsilon_i}$, there exists a point $x^i_t \in \partial B_t$ such that $v_{\epsilon_i}(x^i_t) = \epsilon_i/2$ (cf. Figure 4). By the uniformly Lipschitz continuity of v_{ϵ_i} there exists a $d > 0$ independent of ϵ, such that $B_{d\epsilon_i}(x^i_t) \subset P_i$. Therefore

$$\mathcal{L}^2(P_i) \ge \mathcal{L}^2\Big(\bigcup_{x^i_t \in I^c_{\epsilon_i}} B_{d\epsilon_i}(x^i_t)\Big) \ge rd\epsilon_i/4.$$

On the other hand, since $\Delta v_{\epsilon_i} \to 0$ in B_r,

$$\mathcal{L}^2(P_i)/\epsilon_i \le C \int_{B_r} \beta_{\epsilon_i}(v_{\epsilon_i}) \to 0.$$

Thus we get a contradiction and therefore $\chi_0 \equiv 1$ in B_r. $\square$

Lemma 6.3. *Let* $v_0(x^0) = 0$ *for some* $x^0 \in B_1$, $0 < r_k \to 0$ *as* $k \to \infty$ *and* $v_{0k}(x) = \frac{v_0(x^0 + r_k x)}{r_k}$. *Then there exists a blow-up limit* v_{00} *such that for a subsequence the following holds:*
1) $v_{0k} \to v_{00}$ *in* $C^{0,\alpha}_{\text{loc}}(\mathbb{R}^2)$ *for every* $0 < \alpha < 1$, $\nabla v_{0k} \to \nabla v_{00}$ *weakly* in* $L^\infty_{\text{loc}}(\mathbb{R}^2)$,
2) $v_{0k} \to v_{00}$ *in* $W^{1,2}_{\text{loc}}$,
3) v_{00} *is homogeneous of degree* 1.

Proof. 1) Since $\nabla v_{0k}(x) = \nabla v_0(x_0 + r_k x)$, $\{v_{0k}\}$ is locally uniformly Lipschitz continuous in $\mathbb{R}^2$ and 1) follows.

2) Let $D \subset\subset \mathbb{R}^2$. Then there exists a $C < \infty$ such that $|\nabla v_{0k}(x)| < C$ in D, hence v_{0k} is bounded $W^{1,2}(D)$. It follows that there is a subsequence, which we still call v_{0k}, such that $v_{0k} \to v_{00}$ weakly in $W^{1,2}(D)$ and

$$||v_{00}||_{W^{1,2}(D)} \le \liminf_{k\to\infty} ||v_{0k}||_{W^{1,2}(D)}.$$

Since $\Delta v_0 = \mu$ and *supp* $\mu \subset \partial\{v_0 > 0\}$, we get

$$v_{0k}(x)\Delta v_{0k}(x) = 0.$$

Let $\eta \in C_0^\infty(\mathbb{R}^2)$. Then $\int_{\mathbb{R}^2} \eta |\nabla v_{0k}|^2 = -\int_{\mathbb{R}^2} v_{0k} \nabla\eta \cdot \nabla v_{0k}$ and

$$\limsup_{k\to 0} \int_{\mathbb{R}^2} \eta |\nabla v_{0k}|^2 \le -\int_{\mathbb{R}^2} v_{00} \nabla\eta \cdot \nabla v_{00}.$$

For each $\delta > 0$, we define

$$v_\delta(x) = \begin{cases} v_{00}(x) - \delta, & \text{if } v_{00}(x) > \delta, \\ v_{00}(x) + \delta, & \text{if } v_{00}(x) < -\delta, \\ 0 & \text{otherwise.} \end{cases}$$

Multiplying the equation for v_{0k} by ηv_δ, we get

$$\eta(x) v_\delta(x) \Delta v_{0k}(x) = 0,$$

if k is large enough. Hence

$$\int_{\mathbb{R}^2} \eta \nabla v_\delta \cdot \nabla v_{0k} = -\int_{\mathbb{R}^2} v_\delta \nabla \eta \cdot \nabla v_{0k}.$$

Letting $k \to \infty$, it follows that

$$\int_{\mathbb{R}^2} \eta \nabla v_\delta \cdot \nabla v_{00} = -\int_{\mathbb{R}^2} v_\delta \nabla \eta \cdot \nabla v_{00},$$

and letting $\delta \to 0$, we get

$$\int_{\mathbb{R}^2} \eta |\nabla v_{00}|^2 = -\int_{\mathbb{R}^2} v_{00} \nabla \eta \cdot \nabla v_{00}.$$

Hence

$$\limsup_{k\to 0} \int_{\mathbb{R}^2} \eta |\nabla v_{0k}|^2 \le \int_{\mathbb{R}^2} \eta |\nabla v_{00}|^2$$

for every $\eta \in C_0^\infty(\mathbb{R}^2)$. It follows that $v_{0k} \to v_{00}$ in $W^{1,2}(D)$.

3) For $0 < R < S < \infty$,

$$\Phi(Rr_k) = R^{-2} \int_{B_R} |\nabla v_{0k}|^2 + \chi_0(r_k x)\, dx - R^{-3} \int_{\partial B_R} {v_{0k}}^2 \, d\mathcal{H}^1.$$

Since $v_0(x^0) = 0$ and v_0 is Lipschitz continuous, $\Phi(r)$ is bounded. Consequently we obtain from the monotonicity formula Proposition 5.1 that

$$\begin{aligned} 0 \leftarrow \Phi(Sr_k) - \Phi(Rr_k) &= \int_R^S 2r^{-2} \int_{\partial B_r} (\nabla v_{0k} \cdot \nu - \frac{v_{0k}}{r})^2 d\mathcal{H}^1 dr \\ &= \int_{B_S \setminus B_R} 2|x|^{-4} (\nabla v_{0k}(x) \cdot x - v_{0k}(x))^2 dx. \end{aligned}$$

Letting $k \to \infty$, we obtain that $\nabla v_{00}(x) \cdot x = v_{00}(x)$ a.e. in $\mathbb{R}^2$, hence v_{00} is homogeneous of degree 1. ☐

Now we are ready to prove Theorem A of the Introduction.

Theorem 6.4. *There is a free boundary point $x^0 \in \partial\{v_0 > 0\}$ such that v_0 is degenerate at x^0. More precisely,*

$$\lim_{r\to 0} \frac{v_0(x^0 + rx)}{r} = 0$$

for every $x \in \mathbb{R}^2$.



Figure 5. Touching the Free Boundary

Proof. Case 1: $0 \notin \partial\{v_0 > 0\}$.

There exists an $r > 0$ such that $v_0(x) \leq 0$ in B_r. Since $v_0(0) = 0$, the subharmonicity of v_0 implies that $v_0 \equiv 0$ in B_r. Suppose that r is the largest number such that $v_0 \equiv 0$ in B_r. Noticing that $r < 1$ as the boundary values of v_0 are not constant. It follows that there is a point $x^0 \in \partial B_r$ such that $x^0 \in \partial\{v_0 > 0\}$ (cf. Figure 5).

By a translation and a rotation we may assume that $x^0 = 0$ and $v_0 \equiv 0$ in $B_r(-r, 0)$. Let v_{00} be a blow-up limit of v_0 at x^0, i.e. the limit of $\frac{v_0(x^0 + r_k x)}{r_k}$ as $k \to \infty$. We are going to show that $v_{00} \equiv 0$.

Suppose towards a contradiction that $v_{00} \not\equiv 0$ and let $S = \{v_{00} > 0\}$. Then v_{00} is harmonic in S. Moreover, we know from Lemma 6.3 that v_{00} is homogeneous of degree 1, so that, solving the resulting ODE for v_{00}, $v_{00} = cx \cdot \nu$ in S for some $c \in \mathbb{R}$ and $\nu \in \partial B_1$. Noticing that $v_{00} \equiv 0$ in the left half plane, we conclude that $v_{00} = c \max(0, x \cdot \nu)$, where $c > 0$ and $\nu = (1, 0)$.

Let $\chi_r(x) = \chi_0(rx)$ and let χ_{00} be the limit of χ_{r_k} of the above sequence $k \to \infty$. By Lemma 6.2, $\chi_0 \equiv 1$ in $B_r(-r, 0)$, hence $\chi_{00} \equiv 1$ in the left half plane. Moreover we know that $\chi_0(x) = 1$ if $v_0(x) > 0$, hence $\chi_{00} \equiv 1$ in the right half plane. Thus $\chi_{00}(x) \equiv 1$ in $\mathbb{R}^2$.

From

$$0 = \int_{\mathbb{R}^2} |\nabla v_{00}|^2 \operatorname{div} \phi - 2\nabla v_{00} D\phi \nabla v_{00} + \chi_{00} \operatorname{div} \phi \, dx$$

for every $\phi \in C_0^\infty(\mathbb{R}^2; \mathbb{R}^2)$, we infer that

$$\int_{\mathbb{R}^2} |\nabla v_{00}|^2 \operatorname{div} \phi - 2\nabla v_{00} D\phi \nabla v_{00} = 0$$

for every $\phi \in C_0^\infty(\mathbb{R}^2;\mathbb{R}^2)$. This however contradicts $v_{00} = c\max(0, x\cdot\nu)$. We conclude that the blow-up limit v_{00} must be the constant function 0 in $\mathbb{R}^2$.

Case 2: $0 \in \partial\{v_0 > 0\}$.

Suppose towards a contradiction that a blow-up limit

$$v_{00} = \lim_{j\to\infty} \frac{v_0(s_j\cdot)}{s_j} \not\equiv 0.$$

Similar to case 1, v_{00} must be of the form $cx\cdot\nu$ in some half plane; here ν is again a unit vector. On the other hand Proposition 4.3 5) — which incidentally is preserved under the blow-up limit — implies that there are at least two vectors $e^1 \neq e^2 \in \partial B_1$ and two half lines $\{\alpha e^1 : \alpha > 0\}$ and $\{\alpha e^2 : \alpha > 0\}$, such that $e^1\cdot\nu > 0$, $\nabla v_{00} = \nabla v_{00}\cdot e^1\, e^1$, $e^2\cdot\nu > 0$ and $\nabla v_{00} = \nabla v_{00}\cdot e^2\, e^2$ which is not possible for $v_{00}(x) = cx\cdot\nu$ unless $c = 0$. □

Remark 6.5. Using methods in the forthcoming paper [11], it is actually possible to show that Case 1 in the proof of Theorem 6.4 cannot occur. More precisely, a degenerate point x^0 at which the set $\{v_0 = 0\}$ contains a disk touching x^0 is not possible.

Consequently the origin must in our example be a degenerate point. The fact that we can introduce many symmetry lines suggests that we can construct degenerate points with growth $\leq |x|^m$ in $B_{1/2}$ for any integer m.

REFERENCES

[1] H. W. Alt and L. A. Caffarelli. Existence and regularity for a minimum problem with free boundary. *J. Reine Angew. Math.*, 325:105–144, 1981.

[2] J. Andersson and G. S. Weiss. Cross-shaped and degenerate singularities in an unstable elliptic free boundary problem. *J. Differential Equations*, 228(2):633–640, 2006.

[3] H. Berestycki, L. A. Caffarelli, and L. Nirenberg. Uniform estimates for regularization of free boundary problems. In *Analysis and partial differential equations*, volume 122 of *Lecture Notes in Pure and Appl. Math.*, pages 567–619. Dekker, New York, 1990.

[4] J. D. Buckmaster and G. S. S. Ludford. *Theory of laminar flames*. Cambridge Monographs on Mechanics and Applied Mathematics. Cambridge University

Press, Cambridge, 1982. Electronic & Electrical Engineering Research Studies: Pattern Recognition & Image Processing Series, 2.

[5] Luis A. Caffarelli. Uniform Lipschitz regularity of a singular perturbation problem. *Differential Integral Equations*, 8(7):1585–1590, 1995.

[6] D. Danielli, A. Petrosyan, and H. Shahgholian. A singular perturbation problem for the p-Laplace operator. *Indiana Univ. Math. J.*, 52(2):457–476, 2003.

[7] J. Frehse. Capacity methods in the theory of partial differential equations. *Jahresber. Deutsch. Math.-Verein.*, 84(1):1–44, 1982.

[8] David Gilbarg and Neil S. Trudinger. *Elliptic partial differential equations of second order*, volume 224 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1983.

[9] Alex Gurevich. Boundary regularity for free boundary problems. *Comm. Pure Appl. Math.*, 52(3):363–403, 1999.

[10] Claudia Lederman and Noemi Wolanski. Viscosity solutions and regularity of the free boundary for the limit of an elliptic two phase singular perturbation problem. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 27(2):253–288 (1999), 1998.

[11] E. Varvaruca and G. S. Weiss. A geometric proof of a generalized Stokes conjecture. *In preparation.*

[12] G. S. Weiss. A singular limit arising in combustion theory: fine properties of the free boundary. *Calc. Var. Partial Differential Equations*, 17(3):311–340, 2003.

[13] Georg S. Weiss. Partial regularity for weak solutions of an elliptic free boundary problem. *Comm. Partial Differential Equations*, 23(3-4):439–455, 1998.

Graduate School of Mathematical Sciences, University of Tokyo,
3-8-1 Komaba, Meguro-ku, Tokyo-to, 153-8914 Japan
E-mail address: `gw@ms.u-tokyo.ac.jp`

Graduate School of Mathematical Sciences, University of Tokyo,
3-8-1 Komaba, Meguro-ku, Tokyo-to, 153-8914 Japan
E-mail address: `zhang@ms.u-tokyo.ac.jp`